

Machine Learning

Lecture 2 - Bayesian Learning: Binomial and Dirichlet Distributions

Devdatt Dubhashi
dubhashi@chalmers.se

Department of Computer Science and Engineering
Chalmers University

January 21, 2016

- ▶ Estimate probability a coin shows head based on observed coin tosses.
- ▶ Simple but fundamental!
- ▶ Historically important: originally used by alertBayes (1763) and generalized by Pierre–Simon de Laplace (1774) creating Bayes Rule.

Likelihood

Suppose $X_i \sim \text{Ber}(\theta)$ i.e.

$$P(X_i = 1) = \theta(\text{"heads"}) \quad P(X_i = 0) = 1 - \theta(\text{"tails"})$$

and $\theta \in [0,1]$ is the **parameter** to be estimated.

Given a series of N observed coin tosses, the probability that we observe k heads is given by the

Binomial distribution

:

$$\text{Bin}(k | N, \theta) = \binom{N}{k} \theta^k (1 - \theta)^{N-k}.$$


Thus, the likelihood has the form

$$P(\mathcal{D} | \theta) \propto \theta^{N_1} (1 - \theta)^{N_0},$$

where N_0 and N_1 are the number of tails and heads seen respectively. These are called **sufficient statistics** since this is all we need to know about the data to estimate θ . Formally, $s(\mathcal{D})$ are a set of sufficient statistics for \mathcal{D} if

$$P(\theta | \mathcal{D}) = P(\theta | s(\mathcal{D})).$$

Binomial Distribution



Bayes Rule for Posterior

$$P(\theta | \mathcal{D}) = \frac{P(\mathcal{D} | \theta)P(\theta)}{\int_0^1 P(\mathcal{D} | \theta)P(\theta)d\theta}$$

Bit daunting to compute the integral in the denominator!

Can avoid it via a clever trick of choosing suitable prior!

Prior

Need a prior with support $[0, 1]$ and to make math easy, of the same form as likelihood.

Beta distribution

$$\text{Beta}(\theta \mid a, b) = \frac{1}{B(a, b)} \theta^{a-1} (1 - \theta)^{b-1},$$

with **hyperparameters** a, b , and where

$$B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)},$$

is a normalizing factor.

Mean $\frac{a}{a+b}$


Mode $\frac{a-1}{a+b-2}$

Prior Knowledge If we believe that the mean is 0.7 and standard deviation is 0.2 we can set $a = 2.975$ and $b = 1.277$ (exercise!)

Uninformative Prior Uniform prior $a = 1 = b$.

Beta Distribution

beta distributions



Posterior, Conjugate Prior


Multiplying prior and likelihood gives the posterior:

$$\begin{aligned} P(\theta | \mathcal{D}) &\propto \text{Bin}(N_1 | \theta, N_0 + N_1) \text{Beta}(\theta | a, b) \\ &= \text{Beta}(\theta | N_1 + a, N_0 + b). \end{aligned}$$

The posterior has the same distribution as the prior (with different parameters): the Beta distribution is said to be a **conjugate prior** for the Binomial distribution.

The posterior is obtained by simply adding the prior parameters to the empirical counts, hence the hyperparameters are often called **pseudo-counts**.

Updating Beta Prior with Binomial Likelihood



Sequential update versus Batch

Suppose we have two data sets \mathcal{D}^1 and \mathcal{D}^2 with sufficient statistics N_1^1, N_0^1 and N_1^2, N_0^2 respectively. Let $N_1 := N_1^1 + N_1^2$ and $N_0 := N_0^1 + N_0^2$ be the combined sufficient statistics.

In **batch** mode,

$$\begin{aligned}P(\theta | \mathcal{D}^1, \mathcal{D}^2) &\propto \text{Bin}(N_1 | \theta, N_1 + N_0) \text{Beta}(\theta | a, b) \\ &\propto \text{Beta}(\theta | N_1 + a, N_0 + a).\end{aligned}$$

In **sequential** mode,

$$\begin{aligned}P(\theta | \mathcal{D}^1, \mathcal{D}^2) &\propto P(\mathcal{D}^2 | \theta) P(\theta | \mathcal{D}^1) \\ &\propto \text{Bin}(N_1^2 | \theta, N_1^1 + N_0^2) \text{Beta}(\theta | N_1^1 + a, N_0^1 + b) \\ &\propto \text{Beta}(\theta | N_1^1 + N_1^2 + a, N_0^1 + N_0^2 + b) \\ &= \text{Beta}(\theta | N_1 + a, N_0 + b)\end{aligned}$$

Very suitable for **online learning**!

Posterior Mean and Mode

- ▶ The MAP estimate is given by

$$\hat{\theta}_{MAP} = \frac{a + N_1 - 1}{a + b + N - 2}.$$

- ▶ With uniform prior $a = 1 = b$,

$$\hat{\theta}_{MLE} = \frac{N_1}{N},$$

which is just the MLE.

- ▶ The posterior mean is

$$\bar{\theta} = \frac{a + N_1}{a + b + N},$$

which is a convex combination of the prior mean and the MLE:

$$\bar{\theta} = \lambda \frac{a}{a+b} + (1 - \lambda) \hat{\theta}_{MLE},$$

with $\lambda := \frac{a+b}{a+b+N}$. Note that as $N \rightarrow \infty$, $\bar{\theta} \rightarrow \hat{\theta}_{MLE}$.

The probability of a heads in a single new coin toss is:

$$\begin{aligned}P(\hat{x} = 1 \mid \mathcal{D}) &= \int_0^1 P(x = 1 \mid \theta)P(\theta \mid \mathcal{D})d\theta \\&= \int_0^1 \theta \text{Beta}(\theta \mid N_1 + a, N_0 + b)d\theta \\&= E_{\text{Beta}}[\theta] \\&= \frac{N_1 + a}{N_1 + N_0 + a + b}\end{aligned}$$

Predicting Multiple Future Trials


The probability of predicting x heads in M future trials:

$$\begin{aligned}
 P(x | \mathcal{D}) &= \int_0^1 \text{Bin}(x | \theta, M) P(\theta | \mathcal{D}) d\theta \\
 &= \int_0^1 \text{Bin}(x | \theta, M) \text{Beta}(\theta | N_1 + a, N_0 + b) d\theta \\
 &= \binom{M}{x} \frac{1}{B(N_1 + a, N_0 + b)} \int_0^1 \theta^{x+N_1+a-1} (1-\theta)^{M-x+N_0+b-1} \\
 &= \binom{M}{x} \frac{B(x+N_1+a, (M-x)+N_0+b)}{B(N_1+a, N_0+b)}
 \end{aligned}$$

the **compound Beta–Binomial distribution**, with mean and variance:

$$E[x] = M \frac{N_1 + a}{N + a + b}, \text{var}[x] = M \frac{(N_1 + a)(N_0 + b)}{(N_1 + a + N_0 + b)^2} \frac{N + a + b + M}{N + 1}$$

Posterior Predictive and Plugin



- ▶ From coins and two outcomes to dice and many outcomes.
- ▶ Given observations from a dice with K faces, predict next roll.
- ▶ Suppose we observe N dice rolls $\mathcal{D} = \{x_1, x_2, \dots, x_N\}$ where each $x_i \in \{1, \dots, K\}$.

- ▶ Suppose we observe N dice rolls $\mathcal{D} = \{x_1, x_2, \dots, x_N\}$ where each $x_i \in \{1, \dots, K\}$.



$$P(\mathcal{D} | \theta) = \prod_{k=1}^K \theta_k^{N_k},$$

where θ_k is the (unknown) probability of showing face k and N_k is the observed number outcomes showing face k .

Multinomial Distribution

The probability of observing face k appear x_k times in n rolls of a dice with face probabilities $\theta := (\theta_k, k \in \{1, \dots, K\})$ is

Multinomial Distribution

$$\text{Mu}(\mathbf{x} \mid n, \theta) = \binom{n}{x_1, x_2, \dots, x_K} \prod_{k=1}^K \theta_k^{x_k}.$$

Since the parameter vector θ lives in the K -dimensional simplex

$$S_K := \{(x_1, \dots, x_K) \mid x_1 + \dots + x_K = 1\},$$

we need a prior that

- ▶ has support on this simplex.
- ▶ is ideally also conjugate to the likelihood distribution i.e. multinomial.

Dirichlet Distribution





$$\text{Dir}(\mathbf{x} \mid \alpha) := \frac{1}{B(\alpha)} \prod_{k=1}^K x_k^{\alpha_k - 1} \mathbf{1}[\mathbf{x} \in S_K],$$

where $B(\alpha)$ is the normalization factor

$$B(\alpha) := \frac{\prod_{k=1}^K \Gamma(\alpha_k)}{\Gamma(\alpha_0)},$$

with $\alpha_0 := \sum_{k=1}^K \alpha_k$.

Dirichlet Distribution



Dirichlet Distribution examples

α_0 controls the **strength** i.e. the peak and α_k s control where it occurs:



(1,1,1) Uniform distribution.

(2,2,2) Broad distribution centered at $(1/3, 1/3, 1/3)$

(20,20,20) Narrow distribution centered at $(1/3, 1/3, 1/3)$.

When $\alpha_k < 1$, for all k , we get “spikes” at the corners.

Samples from Dirichlet Distribution



Prior Dirichlet Prior:

$$\text{Dir}(\theta | \alpha) = \frac{1}{B(\alpha)} \prod_{i=1}^K \theta_k^{\alpha_k - 1}$$

Posterior

$$\begin{aligned} P(\theta | \mathcal{D}) &\propto P(\mathcal{D} | \theta)P(\theta) \\ &\propto \prod_{k=1}^K \theta_k^{N_k} \theta_k^{\alpha_k - 1} \\ &= \prod_{k=1}^K \theta_k^{\alpha_k + N_k - 1} \\ &= \text{Dir}(\theta | \alpha_1 + N_1, \dots, \alpha_K + N_K) \end{aligned}$$

MAP estimate using Lagrange Multipliers

$$\begin{aligned} \max_{\theta} \quad & \text{Dir}(\theta \mid \alpha_1 + N_1, \dots, \alpha_K + N_K) \\ &= \prod_{k=1}^K \theta_k^{\alpha_k + N_k - 1} \\ \text{subject to} \quad & \theta_1 + \theta_2 + \dots + \theta_K = 1. \end{aligned}$$

Use **Lagrange multipliers!** Solution:

$$\hat{\theta}_k = \frac{\alpha_k + N_k - 1}{\alpha_0 + N - K}.$$

With uniform prior this becomes $\hat{\theta}_k = N_k/N$.

$$\begin{aligned}P(X = k | \mathcal{D}) &= \int P(X = k | \theta) P(\theta | \mathcal{D}) d\theta \\&= \int P(X = k | \theta_k) [P(\theta_{-k}, \theta_k | \mathcal{D}) d\theta_{-k}] d\theta_k \\&= \int \theta_k P(\theta_k | \mathcal{D}) d\theta_k \\&= E[\theta_k | \mathcal{D}] \\&= \frac{\alpha_k + N_k}{\alpha + N}\end{aligned}$$

Suppose observe the following sentences:

Sentences

Mary had a little lamb, little lamb, little lamb

Mary had a little lamb, it's fleece as white as
snow

Can we predict which word comes next?

Pre-Processing

- ▶ **Vocabulary**: mary, lamb, little, big, fleece, white, black, snow, rain, unk represented by numerals 1 \cdots 10, (where unk stands for “unknown”).
- ▶ Strip away all punctuation and **stop** words like “a”, “as”, “the”
- ▶ **Stemming** i.e. reduce words to base form by removing plurals etc. For example, “running” becomes “run”.
- ▶ This reduces the given sequences to:

1 10, 3 2 3 2 3 2 3 2,

1 10 3 2 10 5 10 6 8.

Bag of Words Model

- ▶ Count **word occurrences** giving for the size 10 vocabulary, the counts:

2, 4, 4, 0, 1, 1, 0, 10, 4.

- ▶ Using $\text{Dir}(1, 1 \dots, 1)$ prior gives:

$$P(X = k | \mathcal{D}) = E[\theta_k | \mathcal{D}] = \frac{\alpha_k + N_k}{\alpha + N} = \frac{1 + N_k}{10 + 17},$$

- ▶ So predictive distribution for next word is:

$(3/27, 5/27, 5/27, 1/27, 2/27, 2/27, 1/27, 2/27, 1/27, 5/27)$

- ▶ whose modes are $X = 2$ (1amb') and $X = 10$ (unk).
- ▶ The words "big", "black", "rain" have non-zero probabilities, even though they have not been seen !