

Variational Inference

Seung-Hoon Na

Chonbuk National University

ELBO (Evidence lower bound)

- Variational lower bound

$$ELBO(q) = -KL(q(\cdot) || \tilde{p}(\cdot, D)) \quad \leftarrow \text{Posterior prob } (p(\mathbf{x}|D)) \text{를 근사}$$

$$ELBO(q) = -KL(q(\cdot) || \tilde{p}(\cdot, \mathbf{x})) \quad \leftarrow p(\mathbf{h}|\mathbf{x}) \text{ prob를 근사}$$

$$\begin{aligned} J(q) &= \sum_{\mathbf{x}} q(\mathbf{x}) \log \frac{q(\mathbf{x})}{\tilde{p}(\mathbf{x})} \\ &= \sum_{\mathbf{x}} q(\mathbf{x}) \log \frac{q(\mathbf{x})}{Z p^*(\mathbf{x})} \\ &= \sum_{\mathbf{x}} q(\mathbf{x}) \log \frac{q(\mathbf{x})}{p^*(\mathbf{x})} - \log Z \\ &= \mathbb{KL}(q || p^*) - \log Z \end{aligned}$$

ELBO



$$L(q) \triangleq -J(q) = -\mathbb{KL}(q || p^*) + \log Z \leq \log Z = \log p(\mathcal{D})$$

ELBO (Evidence lower bound)

$$\begin{aligned}\log p(X) &= \log \int_Z p(X, Z) \\&= \log \int_Z p(X, Z) \frac{q(Z)}{q(Z)} \\&= \log \left(\mathbb{E}_q \left[\frac{p(X, Z)}{q(Z)} \right] \right) \\&\geq \mathbb{E}_q \left[\log \frac{p(X, Z)}{q(Z)} \right] \\&= \mathbb{E}_q [\log p(X, Z)] + H[Z]\end{aligned}$$

← variational lower bound
or ELBO

ELBO (Evidence lower bound)

$$\begin{aligned} KL [q(Z) || p(Z|X)] &= \int_Z q(Z) \log \frac{q(Z)}{p(Z|X)} \\ &= - \int_Z q(Z) \log \frac{p(Z|X)}{q(Z)} \\ &= - \left(\int_Z q(Z) \log \frac{p(X, Z)}{q(Z)} - \int_Z q(Z) \log p(X) \right) \\ &= - \int_Z q(Z) \log \frac{p(X, Z)}{q(Z)} + \log p(X) \int_Z q(Z) \\ &= -L + \log p(X) \end{aligned}$$



variational lower bound

Mean Field Approximation

- Assume the posterior is a fully factorized approximation of the form

$$q(\mathbf{x}) = \prod_i q_i(\mathbf{x}_i)$$

- Solve the optimization problem:

$$\min_{q_1, \dots, q_D} \text{KL}(q || p)$$



$$\log q_j(\mathbf{x}_j) = \mathbb{E}_{-q_j} [\log \tilde{p}(\mathbf{x})] + \text{const}$$

Mean Field Approximation

- E.g.) we have three variables – x_1, x_2, x_3

$$\mathbb{E}_{-q_2} [f(\mathbf{x})] = \sum_{x_1} \sum_{x_3} q(x_1) q_3(x_3) f(x_1, x_2, x_3)$$

- Very similar to Gibbs sampling
- Gibbs sampling: sending **sampled values** between neighboring nodes
- Mean field approximation: send we send **mean values** between nodes

Mean Field Approximation: Derivation

$$L(q) \triangleq -J(q) = \sum_{\mathbf{x}} q(\mathbf{x}) \log \frac{\tilde{p}(\mathbf{x})}{q(\mathbf{x})} \leq \log p(\mathcal{D})$$

$$\begin{aligned} L(q_j) &= \sum_{\mathbf{x}} \prod_i q_i(\mathbf{x}_i) \left[\log \tilde{p}(\mathbf{x}) - \sum_k \log q_k(\mathbf{x}_k) \right] \\ &= \sum_{\mathbf{x}_j} \sum_{\mathbf{x}_{-j}} q_j(\mathbf{x}_j) \prod_{i \neq j} q_i(\mathbf{x}_i) \left[\log \tilde{p}(\mathbf{x}) - \sum_k \log q_k(\mathbf{x}_k) \right] \\ &= \sum_{\mathbf{x}_j} q_j(\mathbf{x}_j) \sum_{\mathbf{x}_{-j}} \prod_{i \neq j} q_i(\mathbf{x}_i) \log \tilde{p}(\mathbf{x}) \\ &\quad - \sum_{\mathbf{x}_j} q_j(\mathbf{x}_j) \sum_{\mathbf{x}_{-j}} \prod_{i \neq j} q_i(\mathbf{x}_i) \left[\sum_{k \neq j} \log q_k(\mathbf{x}_k) + q_j(\mathbf{x}_j) \right] \\ &= \sum_{\mathbf{x}_j} q_j(\mathbf{x}_j) \log f_j(\mathbf{x}_j) - \sum_{\mathbf{x}_j} q_j(\mathbf{x}_j) \log q_j(\mathbf{x}_j) + \text{const} \end{aligned}$$

Mean Field Approximation: Derivation

$$\log f_j(\mathbf{x}_j) \triangleq \sum_{\mathbf{x}_{-j}} \prod_{i \neq j} q_i(\mathbf{x}_i) \log \tilde{p}(\mathbf{x}) = \mathbb{E}_{-q_j} [\log \tilde{p}(\mathbf{x})]$$

- Rewrite $L(q_j)$ by averaging out all the hidden variables except for x_j

$$L(q_j) = -\mathbb{KL}(q_j || f_j)$$

$$q_j(\mathbf{x}_j) = \frac{1}{Z_j} \exp(\mathbb{E}_{-q_j} [\log \tilde{p}(\mathbf{x})])$$

Mean field: Ising model

$$p(\mathbf{x}, \mathbf{y}) = p(\mathbf{x})p(\mathbf{y}|\mathbf{x})$$

- Prior:

$$p(\mathbf{x}) = \frac{1}{Z_0} \exp(-E_0(\mathbf{x}))$$

$$E_0(\mathbf{x}) = - \sum_{i=1}^D \sum_{j \in \text{nbr}_i} W_{ij} x_i x_j$$

- Likelihood:

$$p(\mathbf{y}|\mathbf{x}) = \prod_i p(\mathbf{y}_i|x_i) = \sum_i \exp(-L_i(x_i))$$

Mean field: Ising model

- Posterior:
$$p(\mathbf{x}|\mathbf{y}) = \frac{1}{Z} \exp(-E(\mathbf{x}))$$
$$E(\mathbf{x}) = E_0(\mathbf{x}) - \sum_i L_i(x_i)$$

- The fully factored approximation:

$$q(\mathbf{x}) = \prod_i q(x_i, \mu_i)$$

$$\log \tilde{p}(\mathbf{x}) = x_i \sum_{j \in \text{nbr}_i} W_{ij} x_j + L_i(x_i) + \text{const}$$

➡
$$q_i(x_i) \propto \exp \left(x_i \sum_{j \in \text{nbr}_i} W_{ij} \mu_j + L_i(x_i) \right)$$

Variational Bayes

- Variational Bayes

$$p(\boldsymbol{\theta}|\mathcal{D}) \approx \prod_k q(\boldsymbol{\theta}_k)$$

- Variational Bayes EM
 - To infer both latent variables and parameters

$$p(\boldsymbol{\theta}, \mathbf{z}_{1:N}|\mathcal{D}) \approx q(\boldsymbol{\theta}) \prod_i q_i(\mathbf{z}_i)$$

Variational Bayes: Example - Univariate Gaussian

- Goal: Infer $p(\mu, \lambda | \mathcal{D})$
- The original conjugate prior:

$$p(\mu, \lambda) = \mathcal{N}(\mu | \mu_0, (\kappa_0 \lambda)^{-1}) \text{Ga}(\lambda | a_0, b_0)$$

- But, we use an approximate factored posterior:

$$q(\mu, \lambda) = q_\mu(\mu) q_\lambda(\lambda)$$

Variational Bayes: Example - Univariate Gaussian

- The unnormalized log posterior

$$\begin{aligned}\log \tilde{p}(\mu, \lambda) &= \log p(\mu, \lambda, \mathcal{D}) = \log p(\mathcal{D}|\mu, \lambda) + \log p(\mu|\lambda) + \log p(\lambda) \\ &= \frac{N}{2} \log \lambda - \frac{\lambda}{2} \sum_{i=1}^N (x_i - \mu)^2 - \frac{\kappa_0 \lambda}{2} (\mu - \mu_0)^2 \\ &\quad + \frac{1}{2} \log(\kappa_0 \lambda) + (a_0 - 1) \log \lambda - b_0 \lambda + \text{const}\end{aligned}$$

Variational Bayes: Example - Univariate Gaussian

- Updating $q_\mu(\mu)$

$$\begin{aligned}\log q_\mu(\mu) &= \mathbb{E}_{q_\lambda} [\log p(\mathcal{D}|\mu, \lambda) + \log p(\mu|\lambda)] + \text{const} \\ &= -\frac{\mathbb{E}_{q_\lambda} [\lambda]}{2} \left\{ \kappa_0(\mu - \mu_0)^2 + \sum_{i=1}^N (x_i - \mu)^2 \right\} + \text{const}\end{aligned}$$



$$q_\mu(\mu) = \mathcal{N}(\mu|\mu_N, \kappa_N^{-1})$$

$$\mu_N = \frac{\kappa_0\mu_0 + N\bar{x}}{\kappa_0 + N}, \quad \kappa_N = (\kappa_0 + N)\mathbb{E}_{q_\lambda} [\lambda]$$

Variational Bayes: Example - Univariate Gaussian

- Updating $q_\lambda(\lambda)$

$$\begin{aligned}\log q_\lambda(\lambda) &= \mathbb{E}_{q_\mu} [\log p(\mathcal{D}|\mu, \lambda) + \log p(\mu|\lambda) + \log p(\lambda)] + \text{const} \\ &= (a_0 - 1) \log \lambda - b_0 \lambda + \frac{1}{2} \log \lambda + \frac{N}{2} \log \lambda \\ &\quad - \frac{\lambda}{2} \mathbb{E}_{q_\mu} \left[\kappa_0 (\mu - \mu_0)^2 + \sum_{i=1}^N (x_i - \mu)^2 \right] + \text{const}\end{aligned}$$



$$q_\lambda(\lambda) = \text{Ga}(\lambda|a_N, b_N)$$

$$a_N = a_0 + \frac{N + 1}{2}$$

$$b_N = b_0 + \frac{1}{2} \mathbb{E}_{q_\mu} \left[\kappa_0 (\mu - \mu_0)^2 + \sum_{i=1}^N (x_i - \mu)^2 \right]$$

Variational Bayes EM: Example - Mixtures of Gaussians

$$p(\mathbf{z}, \mathbf{X} | \boldsymbol{\theta}) = \prod_i \prod_k \pi_k^{z_{ik}} \mathcal{N}(\mathbf{x}_i | \boldsymbol{\mu}_k, \boldsymbol{\Lambda}_k^{-1})^{z_{ik}}$$

- Conjugate prior:

$$p(\boldsymbol{\theta}) = \text{Dir}(\boldsymbol{\pi} | \boldsymbol{\alpha}_0) \prod_k \mathcal{N}(\boldsymbol{\mu}_k | \mathbf{m}_0, (\beta_0 \boldsymbol{\Lambda}_k)^{-1}) \text{Wi}(\boldsymbol{\Lambda}_k | \mathbf{L}_0, \nu_0)$$

- Exact posterior: a mixture of K^N distributions
- So, VB approximation:

$$p(\boldsymbol{\theta}, \mathbf{z}_{1:N} | \mathcal{D}) \approx q(\boldsymbol{\theta}) \prod_i q(\mathbf{z}_i)$$

Variational Bayes EM: Example - Mixtures of Gaussians

- Specifically,

$$q(\mathbf{z}, \boldsymbol{\theta}) = q(\mathbf{z}|\boldsymbol{\theta})q(\boldsymbol{\theta}) = \left[\prod_i \text{Cat}(\mathbf{z}_i|\mathbf{r}_i) \right] \left[\text{Dir}(\boldsymbol{\pi}|\boldsymbol{\alpha}) \prod_k \mathcal{N}(\boldsymbol{\mu}_k|\mathbf{m}_k, (\beta_k \boldsymbol{\Lambda}_k)^{-1}) \text{Wi}(\boldsymbol{\Lambda}_k|\mathbf{L}_k, \nu_k) \right]$$

- Derivation of $q(\mathbf{z})$? Variational E-step
- Derivation of $q(\boldsymbol{\theta})$? Variational M-step